

ADDITIVE PROPERTIES OF EVEN PERFECT NUMBERS

YU TSUMURA

ABSTRACT. A positive integer n is said to be perfect if $\sigma(n) = 2n$, where σ denotes the sum of the divisors of n . In this article, we show that if n is an even perfect number, then any integer $m \leq n$ is expressed as a sum of some of divisors of n .

1. INTRODUCTION.

A perfect number is a positive integer whose proper divisors sum up to n itself. Euclid proved that if $2^p - 1$ is prime, then $n = 2^{p-1}(2^p - 1)$ is an even perfect number. Two millennia later, Euler proved the converse.

Theorem 1. *An even positive integer n is perfect if and only if it is of the form $n = 2^{p-1}(2^p - 1)$, where $2^p - 1$ is prime.*

For a proof, see [1, Theorem 1.3.3, p. 22]. So we have a characterization of even perfect numbers. However, it is not known whether there are infinitely many perfect numbers. In addition, we do not know whether there exists an odd perfect number. See [2] for these problems and related unsolved problems.

2. MAIN RESULTS.

If n is a perfect number, then the sum of all its proper divisor is n . Then, what can we say about the sum of some of its proper divisors? Using Theorem 1, we can prove the following theorem.

Theorem 2. *If n is an even perfect number, then any positive integer m less than or equal to n is expressed as a sum of some of divisors of n .*

Before proving it, we give an example. 6 is an even perfect number since its proper divisors 1, 2, and 3 sum up to 6. Theorem 2 says every positive integer $m \leq 6$ can be written as a sum of some of 1, 2, and 3. For $m = 1, 2$, and 3, they are themselves proper divisors, so we do not need to think about them. For $m = 4$, we have $4 = 1 + 3$. For $m = 5$,

we have $5 = 2 + 3$. Finally, since 6 itself is a perfect number, we have $6 = 1 + 2 + 3$.

Let us move on to the proof of Theorem 2

Proof. From now on, any divisor means a proper divisor of n . Suppose n is an even perfect number. By Theorem 1, we can write $n = 2^{p-1}M_p$, where $M_p = 2^p - 1$ is prime. First of all, note that every number m such that $1 \leq m \leq 2^p - 1$ is a sum of some of $1, 2, 2^2, \dots, 2^{p-1}$. (This is just a binary representation of m .) Since $1, 2, 2^2, \dots, 2^{p-1}$ are divisors of n , we could express every number in $S_0 = \{1, 2, 3, \dots, M_p\}$ as a sum of some of divisors. Since we did not use a divisor M_p , we can add it to numbers in S_1 and we see that every number in $S_1 = \{1 + M_p, 2 + M_p, 3 + M_p, \dots, 2M_p\}$ is a sum of some of its divisors. Next, adding $2M_p$ to numbers in S_0 , we see that every number in $S_2 = \{1 + 2M_p, 2 + 2M_p, 3 + 2M_p, \dots, 3M_p\}$ is a sum of some of its divisors. Similarly, adding kM_p with $1 \leq k \leq 2^{p-1} - 1$ to numbers in S_0 , we see that every number in $S_k = \{1 + kM_p, 2 + kM_p, 3 + kM_p, \dots, (k+1)M_p\}$ is a sum of some of its divisors. Since the set of positive integer less than or equal to n is $\bigcup_{k=0}^{2^{p-1}-1} S_k$, we have expressed every m less than or equal to n as a sum of some of divisors of n . \square

The next question is whether this expression is unique or not. The answer is negative. For example, we take a perfect number $n = 6$ and $m = 3$. Then we can express $m = 1 + 2 = 3$ with proper divisors of 6. Hence $m = 3$ can be expressed in two ways. In general, since $M_p = \sum_{i=0}^{p-1} 2^i$, M_p is expressed in two ways. Hence a multiple of M_p is expressed in two ways except for n itself. However, we show that this is only the case in the following theorem.

Theorem 3. *The expression in Theorem 2 is unique except for $m = kM_p$, where $k = 1, 2, 3, \dots, 2^{p-1} - 1$. Also $m = kM_p$ is expressed in exactly two ways for $k = 1, 2, 3, \dots, 2^{p-1} - 1$.*

Proof. First, we have seen that kM_p is expressed in (at least) two ways for $k = 1, 2, 3, \dots, 2^{p-1} - 1$. Now counting multiplicity, there are $2^{2p-1} - 1$ combinations of divisors of n since the number of proper divisors of $n = 2^{p-1}M_p$ is $2p - 1$. (To exclude an empty combination, we subtracted 1 from 2^{2p-1} .) Since $2^{2p-1} - 1 - n = 2^{p-1} - 1$, there are no m other than $m = kM_p$ that is expressed in two ways and kM_p is not expressed in more than two ways. \square

The next natural question is whether the converse is true. That is, if n is an even positive integer and if every positive integer m less than or equal to n is expressed as a sum of some of proper divisors of n , can we

say that n is an even perfect number? Again, the answer is negative. we give a counterexample. Let us consider $n = 20$. Then it is easy to check that 20 is not a perfect number and some of its proper divisors 1, 2, 4, 5, 10 sum up to all positive integers less than or equal to 20.

In conclusion, divisors of an even perfect number n construct not only n itself but also all positive integers less than or equal to n . What a perfect number it is!

REFERENCES

- [1] Richard Crandall and Carl Pomerance. *Prime numbers*. Springer-Verlag, New York, 2001. A computational perspective.
- [2] Richard K. Guy. *Unsolved problems in number theory*. Problem Books in Mathematics. Springer-Verlag, New York, second edition, 1994. Unsolved Problems in Intuitive Mathematics, I.

*Department of Mathematics, Purdue University 150 N. University Street,
West Lafayette, IN 47907-2067
ytsumura@math.purdue.edu*